TECHNICAL INFORMATION BULLETIN
FOR THE USE OF GRANDEVO® BIOINSECTICIDE
ON POME FRUIT FOR MANAGEMENT OF APPLE MAGGOT

FIFRA Section 2(ee) Recommendation
This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

GRANDEVO®
EPA Reg. No. 84059-17

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Insect	Application Method	Product Use Rate per Application	Application Instructions
Pome Fruit Apples, Crabapple, Loquat, Mayhaw, Pears, and Quince	Apple Maggot	Foliar	2 – 3 pounds per acre or 2 – 3 pounds per 100 gallons of water.	Applications should begin as soon as adult flies are active and continue until adult activity is no longer present. Applications should be made on an interval not to exceed 7 days. Grandevo should be part of an IPM program for control of apple maggot that includes rotation or tank-mix with other insecticides labeled for control of apple maggot.
GRANDEVO® has a **pre-harvest interval (PHI)** of **0 days**. Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of **4 hours**.				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

GRANDEVO® is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2014
Marrone Bio Innovations, Inc.
1540 Drew Ave.,
Davis, California 95618
Telephone (530) 750-2800